

Quantifying Microbial Utilization of Petroleum Hydrocarbons in Salt Marsh Sediments by Using the ^{13}C Content of Bacterial rRNA[∇]

Ann Pearson,^{1*} Kimberly S. Kraunz,¹ Alex L. Sessions,² Anne E. Dekas,^{1,2}
William D. Leavitt,¹ and Katrina J. Edwards³

Department of Earth and Planetary Sciences, Harvard University, 20 Oxford St., Cambridge, Massachusetts 02138¹; Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, California 91125²; and Department of Biological Sciences, University of Southern California, Los Angeles, California 90089³

Received 6 May 2007/Accepted 4 December 2007

Natural remediation of oil spills is catalyzed by complex microbial consortia. Here we took a whole-community approach to investigate bacterial incorporation of petroleum hydrocarbons from a simulated oil spill. We utilized the natural difference in carbon isotopic abundance between a salt marsh ecosystem supported by the ^{13}C -enriched C_4 grass *Spartina alterniflora* and ^{13}C -depleted petroleum to monitor changes in the ^{13}C content of biomass. Magnetic bead capture methods for selective recovery of bacterial RNA were used to monitor the ^{13}C content of bacterial biomass during a 2-week experiment. The data show that by the end of the experiment, up to 26% of bacterial biomass was derived from consumption of the freshly spilled oil. The results contrast with the inertness of a nearby relict spill, which occurred in 1969 in West Falmouth, MA. Sequences of 16S rRNA genes from our experimental samples also were consistent with previous reports suggesting the importance of *Gamma*- and *Deltaproteobacteria* and *Firmicutes* in the remineralization of hydrocarbons. The magnetic bead capture approach makes it possible to quantify uptake of petroleum hydrocarbons by microbes in situ. Although employed here at the domain level, RNA capture procedures can be highly specific. The same strategy could be used with genus-level specificity, something which is not currently possible using the ^{13}C content of biomarker lipids.

Coastal environments are threatened by petroleum spills ranging from low-level discharges to catastrophic accidents. Large spills commonly are followed by clean-up efforts, but complete containment is rare. In all cases, remediation ultimately depends on microbial degradation. The rate of this natural bioremediation varies with physical and biological factors (temperature, wind and wave action, macroecology, and microbial community diversity), all of which have been extensively studied and reviewed (3, 4, 23, 27, 42, 69).

The bacterial genera most commonly identified as hydrocarbon degraders include *Pseudomonas*, *Alcanivorax*, *Marinobacter*, and *Cycloclasticus* (gammaproteobacteria) (5, 14, 19, 74); *Sphingomonas* (alphaproteobacteria) (18); gram-positive bacterial genera such as *Staphylococcus* and *Geobacillus* (43, 76); and deltaproteobacterial genera such as *Desulfobacter*, *Desulfococcus*, *Desulfosarcina*, and *Desulfatibacillum* (1, 12, 24, 51, 63). However, because many of the methods for measuring bacterial degradation of petroleum rely on enrichment cultures or isolates (58, 68, 72), the rates at which hydrocarbons are consumed in situ can be difficult to determine. Studies of the degradation of model hydrocarbons by species in pure or enrichment cultures cannot reflect the rates of conversion of oil into biomass by complex microbial communities in their native environments. Since petroleum is a mixture of many compounds with different degradation potentials, estimates based on the uptake of radiolabeled substrates (usually ^{14}C -labeled

n-alkanes) (69) likely do not represent the utilization rate for the total oil. This is especially evident in gas chromatograms of contaminated sediments, where typically an “unresolved complex mixture” (UCM) is a dominant feature in addition to *n*-alkanes (54). Newer methods of stable-isotope labeling (8) or probing (52) combined with molecular analyses of community diversity by PCR-denaturing gradient gel electrophoresis (35, 56) are promising, but they can require long incubation times and the results are difficult to convert to estimates of oil utilization rates. These methods also are subject to the challenges described above, namely, selection of a single molecule as a proxy for a complex, heterogeneous mixture.

To quantify the microbial incorporation of oil in coastal salt marsh habitats, isotope labeling experiments may not be necessary. It is possible to use whole oil and thus avoid many of the concerns raised above. Variations in the natural abundance of the stable isotopes of carbon, ^{12}C and ^{13}C , can be used to trace the carbon sources used by heterotrophic organisms (7, 13). These ratios are expressed as values of $\delta^{13}\text{C}$:

$$\delta^{13}\text{C} = \left(\frac{{}^{13/12}\text{R}_{\text{sample}}}{{}^{13/12}\text{R}_{\text{standard}}} - 1 \right) 1,000$$

Coffin and colleagues (11) demonstrated that the natural ^{13}C abundance of total RNA extracted from a coastal salt marsh could be used to determine the sources of carbon utilized by the active biological community. *Spartina alterniflora*, a common grass found in salt marshes, has a $\delta^{13}\text{C}$ value of approximately -13‰ (48). The values of $\delta^{13}\text{C}$ measured for heterotrophic bacteria living on salt marsh detritus are similar to the values for *Spartina*, as the marsh grass is the primary producer in these ecosystems. In contrast, fossil fuels, which are derived

* Corresponding author. Mailing address: Department of Earth and Planetary Sciences, Harvard University, 20 Oxford St., Cambridge MA 02138. Phone: (617) 384-8392. Fax: (617) 496-4387. E-mail: pearson@eps.harvard.edu.

[∇] Published ahead of print on 14 December 2007.

mainly from marine products, have a lower natural abundance of ^{13}C , resulting in more negative values of $\delta^{13}\text{C}$. Although petroleum mixtures contain a wide variety of hydrocarbon structures, the $\delta^{13}\text{C}$ value of most oils is near -27‰ (55). In this work, our goal was to utilize the difference between the natural isotopic compositions of these substrates (^{13}C -enriched *Spartina* and ^{13}C -depleted petroleum) to monitor uptake of oil-derived carbon into salt marsh bacteria.

We studied whole-community uptake of hydrocarbons using natural-abundance ^{13}C while simultaneously investigating the microbes likely to be responsible for the uptake. The 16S rRNA molecule (rRNA) is a logical "biomarker" that can link isotopic composition and taxonomy (39, 40, 46). Specifically, we combined (i) phylogenetic approaches to obtain sequences of 16S rRNA genes with (ii) carbon isotopic analyses of native rRNA molecules. Since rRNA in actively growing bacteria can account for 20% of the cellular mass (dry weight), it is possible to purify sufficient quantities of total bacterial or group-specific rRNA molecules to subject the material directly to isotopic analysis.

To this end, we used a moving-wire interface that allows ^{13}C analysis of complex, nonvolatile organic samples, such as nucleic acids (9). Recent improvements in the sensitivity of the device, now termed "spooling-wire microcombustion" (SWiM) (15, 60), made this work feasible by providing sample requirements compatible with RNA purification protocols. Typical elemental analyzers require several orders of magnitude more material. The rRNA-SWiM approach is likely to have many future applications in investigations of carbon metabolism in heterogeneous microbial communities.

In September 1969, the oil barge *Florida* spilled 700,000 liters of diesel fuel (no. 2 fuel oil) in Buzzards Bay near West Falmouth, MA (22). This event has become one of the most well-studied oil spills in history. Thirty years after the spill, sediments in nearby Wild Harbor contain remains of degraded oil. The abundance and composition of the residue, known as a UCM, changed only slightly between 1973 and 2000, and the residue remains confined to a narrow 2- to 3-cm horizon now located approximately 10 cm deep in the marsh sediments (54). In the current work, Wild Harbor served as a control. Specifically, we expected to detect no petroleum-related signal in the $\delta^{13}\text{C}$ values of rRNA, since active biodegradation is now undetectable in the contaminated sediment (62). We then used natural-abundance ^{13}C -rRNA-SWiM measurement to quantify the microbial assimilation of hydrocarbons from a modern, simulated spill in a similar environment. We selected nearby Woodneck Marsh, an uncontaminated site, for the simulated spill. This location is a tidally flushed salt marsh similar to Wild Harbor and is also adjacent to Sippewissett Marsh, where community microbial diversity and metabolism have been studied extensively (21, 25, 61). Here we describe both the microbial diversity enriched by addition of no. 2 fuel oil and the use of carbon isotopic signatures of rRNA to quantify incorporation of oil-derived carbon into microbial biomass.

MATERIALS AND METHODS

Wild Harbor samples. A 32-cm-long core was taken from Wild Harbor marsh, West Falmouth, MA, at a temperature of 5°C (early April 2003) using a 10-cm-diameter polyvinyl chloride tube. This core was frozen at -70°C immediately and returned to the laboratory, where it was later cut in situ into 1-cm slices using a

power saw (Ryobi TS230). Each slice was divided into two halves; one half was used for analysis of hydrocarbons by gas chromatography-mass spectrometry, and the other half was used for RNA extraction.

Woodneck Marsh bulk samples. Bulk samples of *Spartina* and wild rose were collected from Woodneck Beach Salt Marsh, Falmouth, MA. In addition, bulk biomass was skimmed from the surface of a shallow tidal pool, and a mussel was collected from the same location. These samples were frozen in $\text{N}_2(l)$ and homogenized with a mortar and pestle. A portion of each bulk sample was prepared for analysis of $\delta^{13}\text{C}$ of total organic carbon using standard sealed-tube combustion procedures (44). Values of $\delta^{13}\text{C}$ were determined using purified CO_2 by dual-inlet isotope ratio mass spectrometry (VG Prism).

Woodneck Marsh incubation. Twelve shallow cores (diameter, 14 cm; depth, 5 cm) were harvested in midsummer and immediately incubated in sterile (baked in air at 450°C for 8 h) Pyrex crystallizing dishes for 0, 1, or 2 weeks under ambient temperature and sunlight conditions; 8 of these cores were incubated with no. 2 fuel oil (Mobil diesel), and 4 were incubated without fuel oil. An excess of no. 2 fuel oil was supplied as an irregular layer of oil-coated sand on top of the eight inoculated samples. The sand was used to keep the oil from forming a continuous film and thus to allow gas exchange. Oil-free zones were observed throughout the experiment, and some areas of the sediment surface remained exposed. To simulate tidal flushing and prevent desiccation, ~ 20 ml of sterile-filtered marsh water was added to the preparations daily. Four of the samples were harvested after 1 week, and the other four were harvested after 2 weeks. The samples were frozen at -80°C for extraction of hydrocarbons and nucleic acids.

Purification of bacterial rRNA. (i) RNA extraction. Total RNA was extracted using methods described previously (46). Some samples were lysed in 50 mM sodium acetate (pH 5.2) containing 0.5% sodium dodecyl sulfate and 0.1 mg/ml proteinase K (Life Technologies), followed by extraction with phenol-chloroform-isoamyl alcohol (25:24:1) and 100% chloroform, and some samples were extracted with RNeasy (Ambion) used according to the manufacturer's instructions. Nucleic acids were precipitated with isopropanol and washed three times with 70% ethanol, and the pellets were resuspended in Nanopure H_2O . Visual inspection of electrophoresis gels confirmed that the total extracts were enriched in RNA but that some samples also contained small amounts of DNA. Humic acids were removed effectively by the RNeasy method in particular, and significant A_{320} generally was not observed for these samples.

(ii) Magnetic bead capture of small-subunit rRNA. Small-subunit rRNA (16S and 23S rRNA) of bacteria was captured from the total RNA extracts using an Ambion MICROExpress bacterial mRNA enrichment kit, with minor modifications to the manufacturer's protocol. The MICROExpress kit contains a probe cocktail designed to bind, capture, and remove bacterial 16S and 23S rRNA molecules to enrich the mRNA in the remaining solution (compatible genera are listed at <http://www.ambion.com/techlib/misc/microbe.html>). Here, the kit was employed to recover the bacterial rRNA fraction from mixed environmental RNA, and the uncaptured (mRNA) fraction was discarded. The total RNA, capture probe cocktail, and binding solution were hybridized and combined with the included paramagnetic beads, as instructed. After capture of the bead-probe-rRNA complex using a magnetic particle separator as described previously (46), the complex was washed and eluted in Nanopure H_2O at 90°C for 60 s. Recovered rRNA was precipitated in 500 μl of isopropanol plus 10 μl of 5 M NaCl, centrifuged (10,000 $\times g$, 15 min), and washed with 70% ethanol.

To remove residual carbon from buffers used in the capture procedure, samples were cleaned using an Ambion MEGAclear RNA purification kit. RNA recovered from the MEGAclear elution step was precipitated in 500 μl of isopropanol plus 10 μl of 5 M NaCl, centrifuged, and washed as described above. Recovered pellets were redissolved in 400 μl of Nanopure H_2O and reprecipitated using high-performance liquid chromatography-grade isopropanol (Fisher). The final pellets were washed three times in 70% ethanol, dried at 60°C , and redissolved in 10 μl of Nanopure H_2O for isotopic analysis. Method blanks were prepared by processing aliquots of Nanopure water that were the same size using the entire hybridization, capture, precipitation, and clean-up procedure.

Isotopic analysis using a SWiM interface. The SWiM system was described by Sessions et al. (60) and is a modification of the original device built by Brand and Dobberstein (9). Briefly, samples in $\sim 1\text{-}\mu\text{l}$ droplets of water are applied to a wire (0.25-mm nickel wire moving at a rate of 0.8 cm/s), which passes sequentially through a cleaning oven (950°C), a drying oven (120°C), and then a combustion oven (750°C) containing wireform CuO. The atmosphere is excluded from the combustion furnace by positive pressure of the helium carrier gas. A portion of the combustion gases flows through a countercurrent Nafion membrane to remove H_2O , through an open split, and then to a ThermoFinnigan 252 isotope ratio mass spectrometer. CO_2 reference gas for calibration of isotopic ratios was supplied via a custom-built interface similar to the ConFlo III device, and samples

were analyzed as sets of five to seven injections spaced at 45-s intervals. The carbon content and $\delta^{13}\text{C}$ values were calculated from the peak areas of m/z 44 and 45 ion chromatograms. The precision of $\delta^{13}\text{C}$ values attainable with this system is roughly 1‰ for samples containing as little as 10 ng C and 0.2‰ for samples containing 100 ng C (60).

Analytical blanks were estimated by repeatedly measuring 1- μl aliquots of sample-free extracts (described above). They typically contained <10% of the typical carbon concentration in processed RNA samples and had $\delta^{13}\text{C}$ values of -23 to -24‰. The contributions to the overall analytical blank arising from the SWiM system are estimated to be <0.2 ng C (55), with the balance of C presumably derived from RNA extraction procedures. Blank contributions were subtracted from all reported sample $\delta^{13}\text{C}$ values, and the error was propagated to the stated uncertainties using equations given by Hayes (26).

Analysis of hydrocarbons. Aliquots of homogenized material were extracted by vortexing wet sediment in a 1:1 mixture of CH_2Cl_2 and H_2O in 50-ml Teflon tubes. The tubes were centrifuged at $5,000 \times g$, and the CH_2Cl_2 layer was moved to a separatory funnel, where it was washed with H_2O . The total organic extract was passed over anhydrous Na_2SO_4 , reacted with HCl-activated Cu to remove elemental sulfur, and fractionated over SiO_2 gel in hexane- CH_2Cl_2 (95:5). Hydrocarbons were analyzed by gas chromatography-mass spectrometry using an Agilent 6890N gas chromatograph interfaced to a 5973 MSD equipped with an HP-5MS column (30 m by 0.25 mm [inside diameter]; film thickness, 0.25 μm). The gas chromatograph temperature program was 65°C for 2 min, increasing to 130°C (0 min) at a rate of 20°C/min, to 280°C (0 min) at a rate of 6°C/min, and finally to 320°C (25 min) at a rate of 3°C/min.

Phylogenetic characterization. Total DNA was extracted using an UltraClean Mega soil DNA kit from MoBio Labs and was concentrated and purified using a Qiagen QIAquick kit. Partial 16S rRNA genes were amplified from Woodneck Marsh samples using universal bacterial primers (0.4 μM each) 27F (5'-AGAG TTTGATCCTGGCTCAG-3') and 1100R (5'-AGGGTTGCGCTCGTTG-3'), a total DNA template (20 ng), and Sigma JumpStart RedTaq Ready Mix for PCR. The thermal cycling conditions were 2 min at 94°C, followed by 30 cycles of 1 min at 94°C, 1 min at 55°C, and 1.5 min at 72°C and then a final extension step of 10 min at 72°C. Duplicates were run for each sample, and 2.5- μl portions of the duplicates were combined in one reaction mixture and amplified for an additional five cycles using a modified "low bias and reconditioning" protocol (67). PCR products were visualized on a 1.0% agarose gel using SYBR green. All reactions were performed using 25- μl (final volume) mixtures. Negative controls (reagents, lab water) were included with every PCR.

PCR products from all samples were cloned using an Invitrogen TOPO TA cloning kit for sequencing with *E. coli* One Shot TOP10 chemically competent cells. Clones were incubated overnight at 37°C on plates containing LB medium with ampicillin (50 $\mu\text{g}/\text{ml}$), inoculated into 5 ml of liquid medium, and incubated for 24 h at 37°C. Plasmids were purified using a Qiagen QIAprep spin miniprep kit. Sequencing was done at the Dana-Farber/Harvard DNA Resource Core (<http://dnaseq.med.harvard.edu/>).

Preliminary 16S rRNA sequence alignments were obtained using BioEdit (<http://www.mbio.ncsu.edu/BioEdit/bioedit.html>) and then were manually inspected for realignment and trimmed to ~700 bp. A check for chimeric sequences suggested that there were no chimeras (Bellerophon server) (30). Maximum likelihood trees were created from the clone sequences using PHYML (20; <http://atgc.lirmm.fr/phyml/>). The substitution model was HKY, and 100 bootstrap replicates were calculated. Sequences from related species and from similar environmental clones were downloaded from the National Center for Biotechnology Information website (<http://www.ncbi.nlm.nih.gov/>) and included in the alignments and trees. *Aquifex aeolicus* served as the outgroup.

Rarefaction curves were calculated using distance-based operational taxonomic unit (OTU) and richness determination (DOTUR) (59). Distance matrices for the DNA sequences of the clones were calculated using default parameters of dnadist in PHYLIP (<http://evolution.genetics.washington.edu/phylip.html>). Sequences were then assigned to OTUs based on the unweighted-pair group method using average linkages (average neighbor) clustering algorithm implemented in DOTUR with default parameters for precision (0.01) and bootstrapping (1,000).

RESULTS

Wild Harbor relict oil spill. The residue of the no. 2 fuel oil spilled from the barge *Florida* was detected at a depth of 8 to 10 cm in the sediment core from Wild Harbor (Fig. 1a). This residue appeared to be a UCM of hydrocarbons, consistent




FIG. 1. (a) Chromatograms of the sediment core from Wild Harbor showing a degraded UCM and no clear signature of *n*-alkanes. (b) Total RNA yield (in μg RNA/g [wet weight] of sediment). (c) $\delta^{13}\text{C}$ values for total RNA (\blacktriangle) and captured bacterial rRNA (\square).

with previous reports (54). The horizons immediately above and below did not contain significant UCM (Fig. 1a).

The RNA yields from this core were low (3.3 to 5.0 μg RNA/g [wet weight]) (Fig. 1b). Assuming that the RNA content of a typical bacterial cell is 6×10^{-14} g and using a conversion based on the estimated growth rate (6), this RNA concentration is consistent with the presence of 2×10^8 to 4×10^8 cells/g of sediment if all of the RNA was from bacterial biomass. This represents an upper boundary for the microbial cell density, as some of the RNA may have been derived from microaerophilic eukaryotes and/or from the roots of the *Spartina* grass itself. The results are also consistent with estimates derived from phospholipid fatty acid concentrations (62). The extractable RNA showed distinct bands representing the 16S and 23S rRNA molecules by gel electrophoresis; this indicated that there were intact ribosomes in the total community and presumably reflected the potential for metabolic activity.

The values of $\delta^{13}\text{C}$ for the total RNA extracts ranged from -15.5 to -14.3‰ (Table 1). The differences between the values for extracts obtained in the region of the UCM and the values for the horizons above and below the residue of the spill were insignificant (Fig. 1c). The values of $\delta^{13}\text{C}$ for the captured fraction of bacterial rRNA were systematically more negative by ~1‰ than the values for the total RNA. Three of the four values were in the range from -16.1 to -15.7‰, while one sample from within the oil spill horizon (9 to 10 cm) had a slightly more positive $\delta^{13}\text{C}$ value than the other samples (-14.9‰). The difference is not significant given the analytical uncertainties, and the direction of the change is opposite what would be expected if the bacteria in this horizon were using oil as a substrate (the $\delta^{13}\text{C}$ for oil is approximately -27‰).

TABLE 1. RNA yields from environmental samples

Sample	RNA concn ($\mu\text{g/g}$) ^a
Wild Harbor core	
7–8 cm	4.3
8–9 cm	3.8
9–10 cm	3.3
10–11 cm	5.0
Woodneck Marsh surface sediment	
Control	55 \pm 13
Incubation with oil for 1 wk	40 \pm 14
Incubation with oil for 2 wk	37 \pm 10

^a The data are data for single measurements for Wild Harbor and for three measurements for Woodneck Marsh.

Woodneck Marsh fuel oil incubation experiment. Eight samples of surface sediment from nearby Woodneck Marsh were amended with no. 2 fuel oil and harvested over a 2-week period. Four additional samples were reserved as controls (no oil). Four of the amended samples were harvested after 1 week, and four were harvested after 2 weeks. The controls were harvested simultaneously with the samples (two at each time point).

(i) Composition and degradation of oil. The control samples contained no detectable hydrocarbon-rich UCM and therefore had an insignificant petroleum component. All of the oil-inoculated samples still contained readily detectable petroleum (Fig. 2a). The chromatograms included a UCM similar to that found in the deeper, Wild Harbor sediments. As a result of the quantity of fresh oil added, *n*-alkanes were also visible in the chromatograms (Fig. 2a). We examined the composition of this oil before and after incubation by using gas chromatography-mass spectrometry. The net uptake of oil represented a very small fraction of the total amount added, and the composition remained unchanged; we thus presumed that the isotopic composition of the residual oil did not differ from the original composition.

(ii) Total RNA. The RNA yields from the Woodneck Marsh samples were 1 order of magnitude higher than those from the Wild Harbor samples (37 to 55 μg RNA/g [wet weight]) (Table 1), indicating that the surface samples had a greater biomass density than the deeper sediments of the Wild Harbor core. The estimates in both cases could have been influenced by a small amount of residual DNA in the extracts (they were not treated with DNase), although visualization on an agarose gel suggested that the majority of the nucleic acid fractions obtained from both Woodneck Marsh and Wild Harbor was RNA. The Woodneck Marsh data suggest that the concentration was approximately 4×10^9 to 6×10^9 cells/g of sediment if all of the RNA was from bacterial biomass. The total RNA content—and therefore probably the total biomass—of the treatment samples decreased relative to the content of the control samples over the duration of the experiment (Fig. 2b), but 16S rRNA sequence data suggested that the total microbial diversities were similar in the two samples (see below).

The values of $\delta^{13}\text{C}$ for the total RNA extracts ranged from -11.5 to -13.0‰ (Table 1). These values also decreased moderately over the course of the experiment (Fig. 2c), and the difference, $\sim 1.5\text{‰}$, was greater than the isotopic heterogene-

ity among the replicate samples (the sample standard deviations were ± 0.2 to 0.4‰). All treatment samples displayed isotopic depletion relative to the control (no-oil) samples.

(iii) Bacterial rRNA. The values of $\delta^{13}\text{C}$ for the captured fraction of bacterial rRNA ranged from -10.3 to -14.7‰ . These values reflect a 4.4‰ difference between the control samples and the oil incubation samples at 2 weeks (Fig. 2c). The results are consistent with the incorporation of oil-derived carbon into bacterial biomass. The 4.4‰ decrease in the value of $\delta^{13}\text{C}$ of captured bacterial rRNA could be used to estimate the fraction of new bacterial biomass (f_{new}) in the samples that was oil derived, either directly (heterotrophically) or indirectly (autotrophic incorporation of respired CO_2).

$$\delta_{\text{new}} = f_{\text{new}}\delta_{\text{oil}} + (1 - f_{\text{new}})\delta_{\text{control}}$$

Using the control bacterial rRNA end member value of $-10.3\text{‰} \pm 0.6\text{‰}$, an assumed composition for oil of $-27.0\text{‰} \pm 0.5\text{‰}$, and the final value of $-14.7\text{‰} \pm 0.5\text{‰}$, the isotopic mass balance suggested that at 2 weeks, $26\% \pm 4\%$ of the carbon in the captured bacterial rRNA originated from the oil.

Woodneck Marsh microbial community. Partial gene sequences of 16S rRNA genes were obtained from three of the 2-week incubation samples and from one of the control samples (also after 2 weeks of incubation). Forty-six clones were sequenced from control sample SM-10, and 138 clones were sequenced from samples SM-3, SM-6, and SM-9 (treatments). The clones sequenced from the control sample included *Alpha*-, *Delta*-, and *Gamma*proteobacteria at the highest frequencies (Fig. 3a). In contrast, the treatment samples contained primarily *Gamma*proteobacteria, *Firmicutes*, *Spirochaetes*, and other or unknown organisms, while the relative number of *Deltaproteobacteria* clones decreased (Fig. 3a). The number of unique OTUs (the OTU threshold was set at 3% divergence)




FIG. 2. (a) Chromatograms of the sediment cores from Woodneck Marsh showing, for incubation with oil, a regular series of *n*-alkanes centered around *n*- C_{20} , which is absent in the control. (b) Total RNA yield (in μg RNA/g [wet weight]) of sediment. (c) $\delta^{13}\text{C}$ values for total RNA (\blacktriangle) and captured bacterial rRNA (\square).




FIG. 3. (a) Frequency of detection of 16S rRNA clones in the control and 2-week treatment samples. Control plot clones were designated SM-10 (46 clones were sequenced), and treatment plot clones were designated SM-3, SM-6, and SM-9 (138 clones were sequenced). (b) Rarefaction curves obtained by DOTUR for the treatment and control samples, using a cutoff value of 97% sequence identity.

detected at similar sampling levels remained approximately the same, regardless of the presence of oil. Calculations using DOTUR (58) yielded 94 unique OTUs among the 138 sequences in the treatment samples (Fig. 3b) and 33 OTUs among the 46 sequences in the control sample (Fig. 3b). These numbers should be interpreted with caution, however, as the numbers of sequences examined were not the same for the two samples (46 sequences versus 138 sequences) and an estimate of community diversity obtained using the set of only 46 control sequences, in particular, was not likely to be very robust. The individual treatment samples had a diversity similar to that

of the control sample; SM-3 contained 36 OTUs, SM-6 contained 43 OTUs, and SM-9 contained 35 OTUs. These numbers are consistent with similar levels of total diversity in all samples.

As expected, the vast majority of clones were closely related to previously reported marine or salt marsh microbial sequences. Many also were closely related to bacteria that grow in limited-oxygen environments, such as *Clostridium*, *Methylococcus*, and *Desulfovibrio* spp. (Fig. 4a, b, and c). Several of the sequences identified in the treatment samples shared 16S rRNA sequence similarity with other clones that are associated




FIG. 4. Maximum likelihood tree of all sequenced *Gammaproteobacteria* (a), *Firmicutes* (b), and *Deltaproteobacteria* (c) clones calculated using PHYML as described in the text. Bootstrap support values are indicated for major nodes having values of >50%. Green, clones from the control samples; orange, clones from the treatment plots. Reference species are indicated by white letters on a black background. No *Firmicutes* were detected among the 45 clones from the control plot.

with hydrocarbon-contaminated environments (29, 49, 75). These clones include clones obtained from gas hydrates, oil wells, and other sediments.

Although *Gammaproteobacteria* comprised about 20% of the clones detected in both the control and treatment samples (Fig. 3a), the distributions differed markedly (Fig. 4a). In the control samples, the clones were related to *Methylococcus*, *Marinobacter*, and *Alkalilimnicola*. Additional groups were detected in the treatment samples. Most of the new clones aligned with *Idiomarina* spp. or *Thiomicrospira* spp. and were similar to other oil-mineralizing consortia. *Idiomarina* spp. are heterotrophic aerobes (31), and *Thiomicrospira* spp. are microaerophilic sulfur oxidizers that fix CO₂ (37).

No *Firmicutes* were identified in the control sample. However, 18% of the clones detected in the treatment samples were related to *Firmicutes*. The species detected were putative anaerobes closely related to *Clostridium* and *Alkaliphilus* (Fig. 4b) and were similar to other clones from gas hydrates, oil wells, and anaerobic hydrocarbon enrichments (53, 75).

Many of the new *Deltaproteobacteria* in the treatment library were similar to clones from hydrocarbon-contaminated systems. These *Deltaproteobacteria* sequences, however, were more evenly distributed among both the control and treatment samples (Fig. 4c). In both types of samples, clones were found that were related to alkane-degrading species, such as "Candidatus *Desulfococcus oleovorans*" Hxd3 (1, 63). Clones whose sequences were 100% identical to the sequence of a clone from an oil-degrading hydrocarbon mat from Etang de Berre (29) were also found in both the treatment and control samples; this group also clustered more closely with *Desulfococcus* spp. than with other sulfate-reducing *Deltaproteobacteria* (Fig. 4c). The preponderance of the remaining clones appeared to be related to sulfate-reducing species of *Deltaproteobacteria* and included *Desulfovibrio* and *Desulfotalea* spp. No clones closely related to *Geobacter* or *Pelobacter* spp. were detected.

Many of the clones in the other major categories (Fig. 3a) also were closely related to clones from hydrocarbon-rich environments, consistent with the overall observation that the experimental clone library reflected many species that have previously been shown to be involved, directly or indirectly, in the metabolism of hydrocarbons (29, 64, 66, 70).

Additional measurements. The SWiM device was used to obtain $\delta^{13}\text{C}$ data for RNA extracts of pure cultures for method development (Table 2). Concurrent analyses of nucleotide standards, bulk RNA standards, and total nucleic acid extracts produced $\delta^{13}\text{C}$ results that had an analytical precision (based on three to five injections) of around $\pm 0.2\text{‰}$ and that were accurate to within $\pm 1.0\text{‰}$ for samples containing 100 nmol of carbon (60). Values of $\delta^{13}\text{C}$ measured for RNA using the SWiM device were consistently 1 to 2‰ enriched in ¹³C relative to the biomass from which they were derived (shown for *E. coli* in Table 2; also measured for *Ferroplasma acidarmanus* [data not shown]). The results agree with previous data which established that the ¹³C content of RNA is influenced by the ribosugar component (7), because sugars tend to be enriched in ¹³C relative to total biomass.

Other carbon isotopic data were obtained for bulk biomass of major components of the Woodneck Marsh salt marsh community. The data include samples of the major primary producer, the C₄ grass *S. alterniflora*, as well as a C₃ plant (wild

TABLE 2. Isotopic compositions of samples from Wild Harbor and Woodneck Marsh

Sample	Type	$\delta^{13}\text{C}$ (‰)	Error (1 σ) (‰)
Wild Harbor ^a			
7.5 cm	Total RNA	-14.6	0.2
8.5 cm	Total RNA	-14.3	0.5
9.5 cm	Total RNA	-14.5	0.5
10.5 cm	Total RNA	-15.5	0.4
7.5 cm	Captured rRNA	-15.8	0.8
8.5 cm	Captured rRNA	-16.1	1.1
9.5 cm	Captured rRNA	-14.9	0.4
10.5 cm	Captured rRNA	-15.7	0.3
Woodneck Marsh ^b			
Control	Total RNA	-11.5	0.2
Oil incubation, 1 wk	Total RNA	-12.0	0.2
Oil incubation, 2 wk	Total RNA	-13.0	0.4
Control	Captured RNA	-10.3	0.6
Oil incubation, 1 wk	Captured rRNA	-12.1	1.3
Oil incubation, 2 wk	Captured rRNA	-14.7	0.5
Bulk samples from Woodneck Marsh			
Grass (<i>S. alterniflora</i>)	Emerged grass	-13.9	0.2 ^c
Wild rose	Whole leaf	-27.4	0.2 ^c
Mussel (<i>G. demissa</i>)	Whole tissue	-14.8	0.2 ^c
Marsh microbial mat total organic carbon	Whole mat & sediment	-15.6	0.2 ^c
Blanks			
Total	Total RNA blank	-23.4	1.2 ^d
Captured	Captured RNA blank	-24.2	1.2 ^d
<i>E. coli</i> controls			
<i>E. coli</i> whole	Whole cells	-22.8	0.2 ^c
<i>E. coli</i> total	Total RNA	-21.5	0.1
<i>E. coli</i> captured	Captured RNA	-20.6	0.6

^a The values of $\delta^{13}\text{C}$ for Wild Harbor are based on triplicate measurements of individual samples.

^b The values of $\delta^{13}\text{C}$ for Woodneck Marsh are based on triplicate measurements for four replicate samples; the errors were calculated by determining sample standard deviations (1 σ).

^c Estimated error for closed-tube combustion.

^d The error for the captured sample was measured; the error for the total sample was assumed to be equivalent to the error for the captured sample.

rose). A marsh mud-dwelling mussel, *Geukensia demissa*, and a mixed sample of whole marsh mud plus bacteria (mat total organic carbon) also were examined.

DISCUSSION

Residual no. 2 fuel oil remains in the sediment in Wild Harbor (54, 71), where it spilled from the barge *Florida* in 1969. Active biodegradation of this residue by the bacterial community is undetectable, as determined previously using natural-abundance ¹⁴C measurements of intact phospholipids extracted from the sediments (62). Our data agree with this previous conclusion; the microbial incorporation of highly de-

graded oil is insignificant. The amount of ^{13}C in bacterial rRNA was uniform, regardless of whether it was purified from samples obtained within or outside the relict, spill-contaminated horizons (Fig. 1c). However, this environment also is different from Woodneck Marsh in several ways that could significantly affect (reduce) the microbial metabolism of hydrocarbons. It is buried deep within the anoxic zone of the marsh sediment and has no contact with oxygenated substrates other than possible irrigation via the *Spartina* roots. The total carbon environment and total microbial community in this anoxic setting also must be different from those at Woodneck Marsh. This is reflected in the $\delta^{13}\text{C}$ value for total RNA in Wild Harbor, which is 3‰ less isotopically (approximately -14.5‰) than the value for untreated total RNA from the surface community in Woodneck Marsh (-11.5‰). This effect could be the result of differences in the carbon substrates and assimilation pathways used by the microbial communities in the two locations.

In the Woodneck Marsh sediments, both the phylogenetic distribution and the ^{13}C content of bacterial rRNA were affected by the addition of fresh fuel oil. The data demonstrate that the salt marsh microbial community responded rapidly to the introduction of hydrocarbon contaminants. The observation that this carbon entered the active metabolic pool—as detected using the labile 16S and 23S rRNA molecules—is consistent with the incorporation of carbon from petroleum into microbial biomass. Biodegradation of hydrocarbons proceeds first through the more labile *n*-alkanes to more recalcitrant molecules, such as aromatic hydrocarbons and branched alkanes (54, 58, 69). The duration of the experiment captured only the initial degradation of the labile fraction of oil, and the experiment was not sensitive enough to detect differential utilization of structurally or isotopically distinct components; a longer experiment is needed to determine changes in the microbial community after the depletion of *n*-alkanes.

However, in this short experiment we estimated that after 2 weeks, up to $\sim 26\%$ of the bacterial carbon was derived from the oil. Stable isotope labeling and analysis of captured rRNA have been used previously to assess microbial uptake of a variety of “tracer” carbon compounds (40). Other studies have examined the catabolic conversion of hydrocarbon substrates into CO_2 (2), while more recent approaches have included isotope labeling methods (DNA stable isotope probing or lipids) to detect the flow of oil-derived carbon into active microbes (8, 41, 47). Many of these studies utilized labeled substrates and/or examined only a single compound. To our knowledge, the study presented here is the first study to use the natural abundance of ^{13}C in rRNA to trace the microbial incorporation of carbon from total oil as a substrate. Conversely, we could not detect if some components of the oil were preferentially consumed relative to other components, other than to note that the total composition of the mixture did not appear to change with time. Regardless, in general, this experiment expanded the potential for using non-isotope-labeled nucleic acids as tracers for natural biogeochemical processes. In the case of petroleum biodegradation, this approach may be particularly well suited.

Oil spills affect sediment-associated microbial consortia by stimulating the growth of species involved directly and indirectly in metabolism of the complex hydrocarbon mixtures (28,

32, 57, 75). In the Woodneck Marsh incubations we observed differences in the community composition between the controls and the treatment experiments, both within groups (*Gammaproteobacteria*) and with the appearance of new groups (*Firmicutes*). The frequency of our clone phylogenies shifted from primarily *Alpha*-, *Gamma*-, and *Deltaproteobacteria* in the control samples to mainly *Gammaproteobacteria*, *Firmicutes*, and *Spirochaetes* in the oil-treated samples. Sulfate-reducing *Deltaproteobacteria* were present in both types of samples, but the relative number of clones decreased and the diversities were comparable in the control and treatment samples. Also, we did not find examples of frequently isolated oil-degrading genera, such as *Azoarcus*, *Cycloclasticus*, or *Alcanivorax*. *Alcanivorax* and *Cycloclasticus* are two of the best-known genera of marine hydrocarbon degraders (14, 33, 34, 56). The failure to detect clones related to these genera may have been due to environmental differences between the Woodneck Marsh salt marsh sediments and other, more open, oceanic sites or could have been due to the relatively short incubation time of the experiment (14 days). The overall biodiversity did not decrease, consistent with some previous reports for hydrocarbon-contaminated marine systems (29, 56).

Many genera of *Gammaproteobacteria* degrade hydrocarbons in pure or enrichment cultures, and clones associated with *Gammaproteobacteria* commonly are detected in environmental incubations and in hydrocarbon-contaminated sediment samples (16, 29, 68, 74, 75). Our enrichment of a set of clones with high similarity to *Idiomarina* spp. (Fig. 4a) also is consistent with the previous detection of this genus in association with oil-rich microbial mats and sediments, (29, 36, 57), although the role that *Idiomarina* plays in the degradation of hydrocarbons is not known.

The sediment microbial community changed visibly when the oil was added, and gradually white filaments appeared to replace pink and purple filaments. This observation is consistent with the results of our clone library sampling analysis; decreased numbers of *Alphaproteobacteria* clone representatives and increased numbers of clones that are related to sulfur oxidizers (among the total clones of *Gammaproteobacteria*) were detected in the library generated for the treatment samples. Many of the new clones observed in the oil-treated samples were associated with *Thiomicrospira* (73) and other sulfur-oxidizing environmental clones (49, 65). Local anoxia also was indicated by the results for the Woodneck Marsh treatment samples. Microbial consortia associated with hydrocarbon degradation in marine sediments frequently contain sulfate reducers (*Deltaproteobacteria*) (49, 57, 66), and sulfate-reducing *Deltaproteobacteria* clones were detected throughout the experiment. Microaerophilic or anoxic conditions also were suggested by the presence of clones affiliated with the genus *Clostridium*, which were absent from the controls.

While we found that the phylogenetic distributions of clones were different for the treatment and control samples, our results do not quantitatively reflect the population changes induced by the mock oil spill. Biases can be introduced by differential copy numbers of 16S rRNA genes or through artifacts of extraction or PCR (10, 17, 38, 50). The results described here thus provide an overview of differences between the oil-treated and control samples. Future work should combine quantitative PCR with the greater phylogenetic specificity of

the ^{13}C -rRNA-SWiM approach in order to obtain a more detailed understanding of the quantitative role that specific microbial groups play in oil degradation.

This work provides a foundation for studies of the metabolic pathways of carbon assimilation in heterogeneous microbial consortia. Here, measurements obtained using the natural isotopic contrast between ^{13}C -depleted oil and a ^{13}C -enriched salt marsh ecosystem showed that hydrocarbons serve as a carbon source at the same time that their oxidation serves as a source of energy. The ^{13}C content of purified bacterial rRNA showed the formation of ^{13}C -depleted biomass over a period of 2 weeks. Most of the isotopically depleted signal was presumed to reflect carbon assimilated from the hydrocarbon amendment, and clonal enrichment of organotrophs is consistent with this observation.

Further study should allow us to investigate hydrocarbon uptake by environmental species using rRNA capture probes having greater specificity. Genus-specific ^{13}C -rRNA-SWiM should be able to quantify the clones detected in the treatment samples, including *Idiomarina* spp., that reflect carbon assimilated directly from oil. We also noted that while the suite of RNA capture probes used here recovers most genera of bacteria, it is not universally compatible. Some excluded organisms of potential environmental interest are *Pirellula* (and presumably other *Planctomycetales*), *Chloroflexus* (and presumably other *Chloroflexales*), and *Thermatogales* (presumably including any of their low-temperature relatives). These exceptions highlight the need for group-targeted probes in future work.

The ^{13}C -rRNA-SWiM approach offers several advantages over traditional substrate incubations. Such approaches provide a valuable new method for studying hydrocarbon assimilation by specific organisms and complement other methods, such as DNA isotope probing (52), labeled-substrate RNA capture (40), and analysis of labeled phospholipid fatty acids (8, 47). Additionally, the investment in technology and labor is arguably less resource-intensive than ion probe (secondary ion mass spectrometry) (45) techniques, although to date our laboratories house the only SWiM interfaces available. The next steps toward broader application of this approach, therefore, include development of new capture probes and broader collaborations between microbiologists and isotope geochemists at the interface of these analytical approaches. Routine $\delta^{13}\text{C}$ analyses of rRNA now require <50 ng C, which is equivalent to the RNA content of 10^4 to 10^6 cells, depending on the growth rate. This should open new windows for understanding carbon flows and pathways in complex microbial consortia.

ACKNOWLEDGMENTS

We are indebted to John M. Hayes for his enduring guidance and support. The manuscript was greatly improved by the detailed comments of two anonymous reviewers. Helen White, Bob Nelson, and Chris Reddy provided access to samples from Wild Harbor. Helen White also gave helpful comments on an early manuscript draft. Sean Sylva, Dan Rogers, Steve Petsch, and Susan Carter are thanked for general laboratory assistance. Al Gagnon provided the analyses of $\delta^{13}\text{C}$ by closed-tube combustion.

This work was supported by a grant from the Reinhart Coastal Research Center of Woods Hole Oceanographic Institution (to A.P. and K.J.E.) and by grant NSF-EAR-0311937 (to A.P.).

REFERENCES

1. Aeckersberg, F., F. Bak, and F. Widdel. 1991. Anaerobic oxidation of saturated hydrocarbons to CO_2 by a new type of sulfate-reducing bacterium. *Arch. Microbiol.* **156**:5–14.
2. Aggarwal, P. K., M. E. Fuller, M. M. Gurgas, J. F. Manning, and M. A. Dillon. 1997. Use of stable oxygen and carbon isotope analyses for monitoring the pathways and rates of intrinsic and enhanced in situ biodegradation. *Environ. Sci. Technol.* **31**:590–596.
3. Atlas, R. M., and R. Bartha. 1973. Stimulated biodegradation of oil slicks using oleophilic fertilizers. *Environ. Sci. Technol.* **7**:538–541.
4. Atlas, R. M. 1981. Microbial degradation of petroleum hydrocarbons—an environmental perspective. *Microbiol. Rev.* **45**:180–209.
5. Bartha, R. 1977. The microbiology of aquatic oil spills. *Adv. Appl. Microbiol.* **22**:225–266.
6. Binder, B. J., and Y. C. Liu. 1998. Growth rate regulation of rRNA content of a marine *Synechococcus* (cyanobacterium) strain. *Appl. Environ. Microbiol.* **64**:3346–3351.
7. Blair, N., A. Leu, E. Muñoz, J. Olsen, E. Kwong, and D. DesMarais. 1985. Carbon isotopic fractionation in heterotrophic microbial metabolism. *Appl. Environ. Microbiol.* **50**:996–1001.
8. Boschker, H. T. S., S. C. Nold, P. Wellsbury, D. Bos, W. deGraaf, R. Pel, R. J. Parkes, and T. E. Cappenberg. 1998. Direct linking of microbial populations to specific biogeochemical processes by C-13-labelling of biomarkers. *Nature* **392**:801–805.
9. Brand, W. A., and P. Dobberstein. 1996. Isotope-ratio-monitoring liquid chromatography mass spectrometry (IRM-LCMS). First results from a moving wire interface system. *Isot. Environ. Health Stud.* **32**:275–283.
10. Chandler, D. P., J. K. Fredrickson, and F. J. Brockman. 1997. Effect of PCR template concentration on the composition and distribution of total community 16S rDNA clone libraries. *Mol. Ecol.* **6**:475–482.
11. Coffin, R. B., D. J. Velinsky, R. Devereux, W. A. Price, and L. A. Cifuentes. 1990. Stable carbon isotope analysis of nucleic acids to trace sources of dissolved substances used by estuarine bacteria. *Appl. Environ. Microbiol.* **56**:2012–2020.
12. Cravo-Laureau, C., R. Matheron, J.-L. Cayol, C. Joulain, and A. Hirschler-Rea. 2004. *Desulfatibacillum aliphaticivorans* gen. nov., sp. nov., an n-alkane- and n-alkene-degrading, sulfate-reducing bacterium. *Int. J. Syst. Evol. Microbiol.* **54**:77–83.
13. DeNiro, M. J., and S. Epstein. 1978. Influence of diet on distribution of carbon isotopes in animals. *Geochim. Cosmochim. Acta* **42**:495–506.
14. Dyksterhouse, S. E., J. P. Gray, R. P. Herwig, J. C. Lara, and J. T. Staley. 1995. *Cycloclasticus pugetii* gen. nov., sp. nov., an aromatic hydrocarbon-degrading bacterium from marine sediments. *Int. J. Syst. Bacteriol.* **45**:116–123.
15. Eek, K. M., A. L. Sessions, and D. P. Lies. 2007. Carbon-isotopic analysis of microbial cells sorted by flow cytometry. *Geobiology* **5**:85–95.
16. Engelhardt, M. A., K. Daly, R. P. Swannell, and I. M. Head. 2001. Isolation and characterization of a novel hydrocarbon-degrading, Gram-positive bacterium, isolated from intertidal beach sediment, and description of *Planococcus alkanolasticus* sp. nov. *J. Appl. Microbiol.* **90**:237–247.
17. Farrelly, V., F. A. Rainey, and E. Stackebrandt. 1995. Effect of genome size and *rrn* gene copy number on PCR amplification of 16S rRNA genes from a mixture of bacterial species. *Appl. Environ. Microbiol.* **61**:2798–2801.
18. Fredrickson, J. K., D. L. Balkwill, G. R. Drake, M. F. Romine, B. Ringelberg, and D. C. White. 1995. Aromatic-degrading *Sphingomonas* isolates from the deep subsurface. *Appl. Environ. Microbiol.* **61**:1917–1922.
19. Gauthier, M. J., B. LaFay, R. Christen, L. Fernandez, M. Acquaviva, P. Bonin, and J. C. Bertrand. 1992. *Marinobacter hydrocarbonoclasticus* gen. nov., sp. nov., a new, extremely halotolerant, hydrocarbon-degrading marine bacterium. *Int. J. Syst. Bacteriol.* **42**:568–576.
20. Gindon, S., and O. Gascuel. 2003. PhyML—a simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Syst. Biol.* **52**:696–704.
21. Hamlett, N. V. 1986. Alteration of a salt-marsh bacterial community by fertilization with sewage sludge. *Appl. Environ. Microbiol.* **52**:915–923.
22. Hampson, G. R., and H. L. Sanders. 1969. Local oil spill. *Oceanus* **15**:8.
23. Harayama, S., Y. Kasai, and A. Hara. 2004. Microbial communities in oil-contaminated seawater. *Curr. Opin. Biotechnol.* **15**:205–214.
24. Harms, G., K. Zengler, R. Rabus, F. Aeckersberg, D. Minz, R. Rosselló-Mora, and F. Widdel. 1999. Anaerobic oxidation of *o*-xylene, *m*-xylene, and homologous alkylbenzenes by new types of sulfate-reducing bacteria. *Appl. Environ. Microbiol.* **65**:999–1004.
25. Harvey, R. W., and L. Y. Young. 1980. Enrichment and association of bacteria and particulates in salt marsh surface water. *Appl. Environ. Microbiol.* **39**:894–899.
26. Hayes, J. M. 1983. Practice and principles of isotopic measurements in organic geochemistry, p. 5.1–5.31. *In* W. G. Meinschein (ed.), *Organic geochemistry of contemporaneous and ancient sediments*. Great Lakes Section, Society of Economic Paleontologists and Mineralogists, Bloomington, IN.
27. Head, I. M., and R. P. J. Swannell. 1999. Bioremediation of petroleum

- hydrocarbon contaminants in marine habitats. *Curr. Opin. Biotechnol.* **10**: 234–239.
28. Hedlund, B. P., A. D. Geiselbrecht, T. J. Bair, and J. T. Staley. 1999. Polycyclic aromatic hydrocarbon degradation by a new marine bacterium, *Neptunomonas naphthovorans* gen. nov., sp. nov. *Appl. Environ. Microbiol.* **65**:251–259.
 29. Hernandez-Raquet, G., H. Budzinski, P. Caumette, P. Dabert, K. Le Menach, G. Muyzer, and R. Duran. 2006. Molecular diversity studies of bacterial communities of oil polluted microbial mats from the Etang de Berre (France). *FEMS Microbiol. Ecol.* **58**:550–562.
 30. Huber, T., G. Faulkner, and P. Hugenoltz. 2004. Bellerophon; a program to detect chimeric sequences in multiple sequence alignments. *Bioinformatics* **20**:2317–2319.
 31. Ivanova, E. P., L. A. Romanenko, J. Chun, M. H. Matte, G. R. Matte, V. V. Mikhailov, V. I. Svetashev, A. Hug, T. Maugel, and R. R. Colwell. 2000. *Idiomarina* gen. nov., comprising novel indigenous deep-sea bacteria from the Pacific Ocean, including descriptions of two species, *Idiomarina abyssalis* sp. nov. and *Idiomarina zobellii* sp. nov. *Int. J. Syst. Evol. Microbiol.* **50**:901–907.
 32. Kasai, Y., H. Kishira, K. Sytsubo, and S. Harayama. 2001. Molecular detection of marine bacterial populations on beaches contaminated by the Nakhodka tanker oil-spill accident. *Environ. Microbiol.* **3**:246–255.
 33. Kasai, Y., H. Kishira, and S. Harayama. 2002. Bacteria belonging to the genus *Cycloclasticus* play a primary role in the degradation of aromatic hydrocarbons released in a marine environment. *Appl. Environ. Microbiol.* **68**:5625–5633.
 34. Kasai, Y., H. Kishira, T. Sasaki, K. Sytsubo, K. Watanabe, and S. Harayama. 2002. Predominant growth of *Alcanivorax* strains in oil-contaminated and nutrient-supplemented sea water. *Environ. Microbiol.* **4**:141–147.
 35. Kasai, Y., Y. Takahata, M. Manefield, and K. Watanabe. 2006. RNA-based stable isotope probing and isolation of anaerobic benzene-degrading bacteria from gasoline-contaminated groundwater. *Appl. Environ. Microbiol.* **72**: 3586–3592.
 36. Kleinstuber, S., V. Riis, I. Fetzer, H. Harms, and S. Muller. 2006. Population dynamics within a microbial consortium during growth on diesel fuel in saline environments. *Appl. Environ. Microbiol.* **72**:3531–3542.
 37. Kuenen, J. G., and R. F. Beudeker. 1982. Microbiology of thiobacilli and other sulphur-oxidizing autotrophs, mixotrophs and heterotrophs. *Phil. Trans. R. Soc. London* **298**:473–497.
 38. Luna, G. M., A. Dell'Anno, and R. Danovaro. 2006. DNA extraction procedure: a critical issue for bacterial diversity assessment in marine sediments. *Environ. Microbiol.* **8**:308–320.
 39. MacGregor, B. J., V. Bruchert, S. Fleischer, and R. Amann. 2002. Isolation of small-subunit rRNA for stable isotopic characterization. *Environ. Microbiol.* **4**:451–464.
 40. MacGregor, B. J., H. T. S. Boschker, and R. Amann. 2006. Comparison of rRNA and polar-lipid-derived fatty acid biomarkers for assessment of ¹³C-substrate incorporation by microorganisms in marine sediments. *Appl. Environ. Microbiol.* **72**:5246–5253.
 41. Madsen, E. L. 2006. The use of stable isotope probing techniques in bioreactor and field studies on bioremediation. *Curr. Opin. Biotechnol.* **17**:92–97.
 42. Margesin, R., M. Hammerle, and D. Tschirko. 2007. Microbial activity and community composition during bioremediation of diesel-oil-contaminated soil: effects of hydrocarbon concentration, fertilizers, and incubation time. *Microb. Ecol.* **53**:259–269.
 43. Maugeri, T. L., C. Gugliandolo, D. Caccamo, and E. Stackebrandt. 2002. Three novel halotolerant and thermophilic *Geobacillus* strains from shallow marine vents. *Syst. Appl. Microbiol.* **25**:450–455.
 44. McNichol, A. P., E. A. Osborne, A. R. Gagnon, B. Fry, and G. A. Jones. 1994. TIC, TOC, DIC, DOC, PIC, POC—unique aspects in the preparation of oceanographic samples for ¹⁴C-AMS. *Nucl. Instr. Methods Phys. Res. B* **92**:162–165.
 45. Orphan, V. J., C. H. House, K. U. Hinrichs, K. D. McKeegan, and E. F. DeLong. 2001. Coupled isotopic and phylogenetic characterization of single cells: direct evidence for a methane-consuming archaeal/bacterial consortium. *Science* **293**:484–487.
 46. Pearson, A., A. L. Sessions, K. J. Edwards, and J. M. Hayes. 2004. Phylogenetically specific separation of rRNA from prokaryotes for isotopic analysis. *Mar. Chem.* **92**:295–306.
 47. Pelz, O., N. Chatzinotas, N. Andersen, S. M. Bernasconi, C. Hesse, J. Abraham, and W.-R. Zeyer. 2001. Use of isotopic and molecular techniques to link toluene degradation in denitrifying aquifer microcosms to specific microbial populations. *Arch. Microbiol.* **175**:270–281.
 48. Peterson, B. J., R. W. Howarth, and R. H. Garritt. 1985. Multiple stable isotopes used to trace the flow of organic matter in estuarine food webs. *Science* **227**:1361–1363.
 49. Phelps, C. D., L. J. Kerkhof, and L. Y. Young. 1998. Molecular characterization of a sulfate-reducing consortium which mineralizes benzene. *FEMS Microbiol. Ecol.* **27**:269–279.
 50. Polz, M. F., and C. M. Cavanaugh. 1998. Bias in template-to-product ratios in multitemplate PCR. *Appl. Environ. Microbiol.* **64**:3724–3730.
 51. Rabus, R., R. Nordhaus, W. Ludwig, and F. Widdel. 1993. Complete oxidation of toluene under strictly anoxic conditions by a new sulfate-reducing bacterium. *Appl. Environ. Microbiol.* **59**:1444–1451.
 52. Radajewski, S., P. Ineson, N. R. Parekh, and J. C. Murrell. 2000. Stable-isotope probing as a tool in microbial ecology. *Nature* **403**:646–649.
 53. Ravot, G., M. Magot, M. L. Fardeau, B. K. Patel, P. Thomas, J. L. Garcia, and B. Ollivier. 1999. *Fusibacter paucivorans* gen. nov., sp. nov., an anaerobic, thiosulfate-reducing bacterium from an oil-producing well. *Int. J. Syst. Bacteriol.* **49**:1141–1147.
 54. Reddy, C. M., T. I. Eglinton, A. Hounshell, H. K. White, L. Xu, R. B. Gaines, and G. S. Frysinger. 2002. The West Falmouth oil spill after thirty years: the persistence of petroleum hydrocarbons in marsh sediments. *Environ. Sci. Technol.* **36**:4754–4760.
 55. Reddy, C. M., L. Xu, J. G. Quinn, and P. C. Hartmann. 2002. Investigating the radiocarbon content of the unresolved complex mixture. *Abstr. Am. Chem. Soc. Div. Environ. Chem.* **224**:32.
 56. Röling, W. F. M., M. G. Milner, D. M. Jones, K. Lee, F. Daniel, R. J. Swannell, and I. M. Head. 2002. Robust hydrocarbon degradation and dynamics of bacterial communities during nutrient-enhanced oil spill bioremediation. *Appl. Environ. Microbiol.* **68**:5537–5548.
 57. Röling, W. F. M., M. G. Milner, D. M. Jones, F. Fratapietro, R. P. Swannell, F. Daniel, and I. M. Head. 2004. Bacterial community dynamics and hydrocarbon degradation during a field-scale evaluation of bioremediation on a mudflat beach contaminated with buried oil. *Appl. Environ. Microbiol.* **70**: 2603–2613.
 58. Rueter, P., R. Rabus, H. Wilkes, F. Aeckersberg, F. A. Rainey, H. W. Jannasch, and F. Widdel. 1994. Anaerobic oxidation of hydrocarbons in crude oil by new types of sulphate-reducing bacteria. *Nature* **372**:455–458.
 59. Schloss, P. D., and J. Handelsman. 2005. Introducing DOTUR, a computer program for defining operational taxonomic units and estimating species richness. *Appl. Environ. Microbiol.* **71**:1501–1506.
 60. Sessions, A. L., S. P. Sylva, and J. M. Hayes. 2005. Moving-wire device for carbon isotopic analyses of nanogram quantities of nonvolatile organic carbon. *Anal. Chem.* **77**:6519–6527.
 61. Simmons, S. L., and K. J. Edwards. 2007. Unexpected diversity in populations of the many-celled magnetotactic prokaryote. *Environ. Microbiol.* **9**:206–215.
 62. Slater, G. F., H. K. White, T. I. Eglinton, and C. M. Reddy. 2005. Determination of microbial carbon sources in petroleum contaminated sediments using molecular C-14 analysis. *Environ. Sci. Technol.* **39**:2552–2558.
 63. So, C. M., and L. Y. Young. 1999. Isolation and characterization of a sulfate-reducing bacterium that anaerobically degrades alkanes. *Appl. Environ. Microbiol.* **65**:2969–2976.
 64. Suzuki, Y., F. Inagaki, K. Takai, K. H. Nealson, and K. Horikoshi. 2004. Microbial diversity in inactive chimney structures from deep-sea hydrothermal systems. *Microb. Ecol.* **47**:186–196.
 65. Takai, K., H. Hirayama, T. Nakagawa, Y. Suzuki, K. H. Nealson, and K. Horikoshi. 2004. *Thiomicrospira thermophila* sp. nov., a novel microaerobic, thermotolerant, sulfur-oxidizing chemolithomixotroph isolated from a deep-sea hydrothermal fumarole in the TOTO caldera, Mariana Arc, Western Pacific. *Int. J. Syst. Evol. Microbiol.* **54**:2325–2333.
 66. Teske, A., K.-U. Hinrichs, V. Edgcomb, A. D. Gomez, D. Kysela, S. P. Sylva, M. L. Sogin, and H. W. Jannasch. 2002. Microbial diversity of hydrothermal sediments in the Guaymas Basin: evidence for anaerobic methanotrophic communities. *Appl. Environ. Microbiol.* **68**:1994–2007.
 67. Thompson, J. R., L. A. Marcelino, and M. F. Polz. 2002. Heteroduplexes in mixed-template amplifications: formation, consequence and elimination by 'reconditioning PCR.' *Nucleic Acids Res.* **30**:2083–2088.
 68. Wang, Y., P. C. Lau, and D. K. Button. 1996. A marine oligobacterium harboring genes known to be part of aromatic hydrocarbon degradation pathways of soil pseudomonads. *Appl. Environ. Microbiol.* **62**:2169–2173.
 69. Ward, D. W., R. M. Atlas, P. D. Boehm, and J. A. Calder. 1980. Microbial biodegradation and chemical evolution of oil from the Amoco spill. *AMBIO* **9**:277–283.
 70. Werne, J. P., R. R. Haese, T. Zitter, G. Aloisi, I. Bouloubassi, S. Heijs, A. Fiala-Medioni, R. D. Pancost, J. S. Sinningh-Damste, G. de Lange L. J. Forney, J. C. Gottschal, J.-P. Foucher, J. Masche, and J. Woodside. 2004. Life at cold seeps: a synthesis of biogeochemical and ecological data from Kazan mud volcano, eastern Mediterranean Sea. *Chem. Geol.* **205**:367–390.
 71. White, H. K., C. M. Reddy, and T. I. Eglinton. 2005. Isotopic constraints on the fate of petroleum residues sequestered in salt marsh sediments. *Environ. Sci. Technol.* **39**:2545–2551.
 72. Whyte, L. G., L. Bourbonniere, and C. W. Greer. 1997. Biodegradation of petroleum hydrocarbons by psychrotrophic *Pseudomonas* strains possessing both alkane (*alk*) and naphthalene (*nah*) catabolic pathways. *Appl. Environ. Microbiol.* **63**:3719–3723.
 73. Wirsén, C. O., T. Brinkhoff, J. Kuever, G. Muyzer, S. Molyneaux, and H. W. Jannasch. 1998. Comparison of a new *Thiomicrospira* strain from the mid-

- Atlantic ridge with known hydrothermal vent isolates. *Appl. Environ. Microbiol.* **64**:4057–4059.
74. **Yakimov, M. M., P. N. Golyshin, S. Lang, E. R. Moore, W. R. Abraham, H. Lunsdorf, and K. N. Timmis.** 1998. *Alcanivorax borkumensis* gen. nov., sp. nov., a new, hydrocarbon-degrading and surfactant-producing marine bacterium. *Int. J. Syst. Bacteriol.* **48**:339–348.
75. **Yakimov, M. M., R. Denaro, M. Genovese, S. Cappello, G. D'Auria, T. N. Chernikova, K. N. Timmis, P. N. Golyshin, and L. Giluliano.** 2005. Natural microbial diversity in superficial sediments of Milazzo Harbor (Sicily) and community successions during microcosm enrichment with various hydrocarbons. *Environ. Microbiol.* **7**:1426–1441.
76. **Zhuang, W. Q., J. H. Tay, A. M. Maszenan, and S. T. Tay.** 2003. Isolation of naphthalene-degrading bacteria from tropical marine sediments. *Water Sci. Technol.* **47**:303–308.